

Vanishing dimension five proton decay operators in the $SU(5)$ SUSY GUT

Naoyuki Haba^{1,2} and Toshihiko Ota¹

¹*Physik Department, Technische Universität München, James-Frank-Strasse 85748 Garching, Germany*

²*Institute of Theoretical Physics, University of Tokushima, 770-8502 Tokushima, Japan*

(Dated: February 2, 2008)

We propose a natural framework of $SU(5)$ supersymmetric grand unified theory with the minimal particle contents, which does not contain dimension five proton decay operators. The suitable fermion mass hierarchy can be reproduced by higher dimensional operators of an adjoint Higgs field which breaks $SU(5)$ gauge symmetry.

PACS numbers: 12.10.Dm, 12.60.Jv, 13.30.-a, 14.20.Dh

Supersymmetric (SUSY) grand unified theories (GUTs)[1] have been regarded as the most agreeable candidates beyond the standard model for a long time, because they can realize the gauge coupling unification as well as show a natural solution of the hierarchy problem. They can also explain the electroweak symmetry breaking by the so-called the radiative breaking scenario[2]. The proton decay is a crucial prediction of GUTs[3], which has not been observed in the experiments yet[4]. From the current experimental bound on the proton decay, it was claimed that the minimal $SU(5)$ SUSY GUT has already been excluded[5, 6]. However, as pointed out in e.g. Refs.[7, 8], this claim should only be applied to the minimal scenario which is unrealistic in the sense that it cannot reproduce the correct mass spectrum of down type quarks and charged leptons. Notice that we must analyze the Yukawa interactions with the coloured Higgs fields carefully. In this letter we show the natural model which does not contain dimension five proton decay operators in the $SU(5)$ SUSY GUT framework with the minimal particle content. We must introduce the GUT scale (M_{GUT}) where the $SU(5)$ gauge symmetry is broken by the vacuum expectation value (VEV) of an adjoint representation (**24**) Higgs field Σ . Since the model at the GUT scale is the effective theory of the fundamental one which realized at the Planck scale, it should contain the higher dimensional operators which are suppressed by the Planck scale ($M_{\text{GUT}}/M_{\text{Pl}}$) ^{n} (n : positive integer) [8, 9, 10]. These terms can be the origin of a fermion mass hierarchy of three generations as well as the one between top and bottom quarks. The realistic mass spectrum of the down type quarks and the charged leptons can be reproduced by these terms. We take the following setup;

1. Only the top Yukawa coupling exists at the tree level. The other Yukawa couplings are induced by the n th order higher dimensional terms, ($M_{\text{GUT}}/M_{\text{Pl}}$) ^{n} .
2. The Yukawa couplings of the bottom quark and the tau lepton are induced by the *first order* terms ($M_{\text{GUT}}/M_{\text{Pl}}$), the mass of the strange quark and the muon are reproduced by the *second order* terms

($M_{\text{GUT}}/M_{\text{Pl}}$)², and the down quark and the electron masses are provided by the *third order* terms ($M_{\text{GUT}}/M_{\text{Pl}}$)³. The masses of the up type quarks might be appropriately reproduced in a similar way.

3. We should pay attention that some of the terms ($\langle \Sigma \rangle / M_{\text{Pl}}$) ^{n} are regarded as $(n-1)$ th order terms effectively due to their coefficients.
4. The couplings of the operators associated with the proton decay process completely vanish.

As we will show, the two requirements — the realistic Yukawa couplings of quarks and leptons and no dimension five proton decay operators — almost determine the couplings for the higher dimensional operators.

The superpotential for the Yukawa sector is represented as the series expansion according to the power of adjoint Higgs fields, such as

$$W_{\text{Yukawa}} = W_0 + W_1 + W_2 + W_3 + W_4 + \cdots \quad (1)$$

The zeroth order part W_0 is the same as that of the minimal $SU(5)$ model,

$$W_0 = \frac{1}{4} \epsilon_{abcde} Y_1^{ij} \mathbf{10}_i^{ab} \mathbf{10}_j^{cd} H^e + \sqrt{2} Y_2^{ij} \bar{H}_a \mathbf{10}_i^{ab} \mathbf{5}_{jb}^*, \quad (2)$$

where i, j are the indices for generations and $a, b, \dots$ are the ones for the $SU(5)$ indices. The chiral superfield $\mathbf{10}$ contains the right-handed up-type quark u_R^c , the left-handed quark doublet Q , and the right-handed charged lepton e_R^c . The right-handed down-type quark d_R^c and the lepton doublet L belong to the superfield $\mathbf{5}^*$. The Higgs fields H and $\bar{H}$ (**5** and **5**^{*}) include the coloured Higgs triplets ($H_C, \bar{H}_C$) and the Higgs doublets, respectively. As shown in our setup, $Y_2 = 0$ is assumed in Eq.(2), which is the origin of the hierarchy between the top and the bottom masses. The *first order* part W_1 is expressed as[8]

$$W_1 = \frac{\epsilon_{abcde}}{4} \left(f_1^{ij} \mathbf{10}_i^{ab} \mathbf{10}_j^{cd} \frac{\Sigma_f^e}{M_{\text{Pl}}} H^f + f_2^{ij} \mathbf{10}_i^{ab} \mathbf{10}_j^{cf} H^d \frac{\Sigma_f^e}{M_{\text{Pl}}} \right) + \sqrt{2} \left(h_1^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \mathbf{5}_{jc}^* + h_2^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{\Sigma_b^c}{M_{\text{Pl}}} \mathbf{5}_{jc}^* \right), \quad (3)$$

where Σ takes VEV of $\langle \Sigma \rangle = \text{diag}(2, 2, 2, -3, -3)\sigma$ which breaks the $SU(5)$ gauge group into $SU(3)_c \times SU(2)_L \times$

$U(1)_Y$. The value of σ is the scale of M_{GUT} . For the *second* and *third order* superpotential, we only show the down-type quark and the charged lepton sector. They are represented as[9]

$$\begin{aligned}
W_2 &= \sqrt{2} \left(h_3^{ij} \bar{H}_a \mathbf{10}_i^{ab} \mathbf{5}_{jb}^* \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} + h_4^{ij} \bar{H}_a \frac{(\Sigma\Sigma)_b^a}{M_{\text{Pl}}^2} \mathbf{10}_i^{bc} \mathbf{5}_{jc}^* \right. \\
&\quad \left. + h_5^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{(\Sigma\Sigma)_b^c}{M_{\text{Pl}}^2} \mathbf{5}_{jc}^* + h_6^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \frac{\Sigma_c^d}{M_{\text{Pl}}} \mathbf{5}_{jd}^* \right), \quad (4) \\
W_3 &= \sqrt{2} \left(h_7^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \mathbf{5}_{jc}^* \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} \right. \\
&\quad + h_8^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{\Sigma_b^c}{M_{\text{Pl}}} \mathbf{5}_{jc}^* \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} + h_9^{ij} \bar{H}_a \mathbf{10}_i^{ab} \mathbf{5}_{jb}^* \frac{(\Sigma\Sigma\Sigma)}{M_{\text{Pl}}^3} \\
&\quad + h_{10}^{ij} \bar{H}_a \frac{(\Sigma\Sigma\Sigma)_b^a}{M_{\text{Pl}}^3} \mathbf{10}_i^{bc} \mathbf{5}_{jc}^* + h_{11}^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{(\Sigma\Sigma\Sigma)_b^c}{M_{\text{Pl}}^3} \mathbf{5}_{jc}^* \\
&\quad \left. + h_{12}^{ij} \bar{H}_a \frac{(\Sigma\Sigma)_b^a}{M_{\text{Pl}}^2} \mathbf{10}_i^{bc} \frac{\Sigma_c^d}{M_{\text{Pl}}} \mathbf{5}_{jd}^* + h_{13}^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \frac{(\Sigma\Sigma)_c^d}{M_{\text{Pl}}^2} \mathbf{5}_{jd}^* \right), \quad (5)
\end{aligned}$$

where $(\Sigma \dots)$ ($(\Sigma \dots)_b^a$) denotes a singlet (an adjoint) by contracting the $SU(5)$ indices of $\Sigma \dots$. The *fourth order* superpotential suggests

$$\begin{aligned}
W_4 &= \sqrt{2} \left(h_{14}^{ij} \bar{H}_a \mathbf{10}_i^{ab} \mathbf{5}_b^* \frac{(\Sigma\Sigma)(\Sigma\Sigma)}{M_{\text{Pl}}^4} \right. \\
&\quad + h_{15}^{ij} \bar{H}_a \mathbf{10}_i^{ab} \mathbf{5}_b^* \frac{(\Sigma\Sigma\Sigma\Sigma)}{M_{\text{Pl}}^4} + h_{16}^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \mathbf{5}_c^* \frac{(\Sigma\Sigma\Sigma)}{M_{\text{Pl}}^3} \\
&\quad + h_{17}^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{\Sigma_b^c}{M_{\text{Pl}}} \mathbf{5}_c^* \frac{(\Sigma\Sigma\Sigma)}{M_{\text{Pl}}^3} + h_{18}^{ij} \bar{H}_a \frac{(\Sigma\Sigma)_b^a}{M_{\text{Pl}}^2} \mathbf{10}_i^{bc} \mathbf{5}_c^* \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} \\
&\quad + h_{19}^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{(\Sigma\Sigma)_b^c}{M_{\text{Pl}}^2} \mathbf{5}_c^* \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} + h_{20}^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \frac{\Sigma_c^d}{M_{\text{Pl}}} \mathbf{5}_d^* \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} \\
&\quad + h_{21}^{ij} \bar{H}_a \frac{(\Sigma\Sigma\Sigma)_b^a}{M_{\text{Pl}}^4} \mathbf{10}_i^{bc} \mathbf{5}_c^* + h_{22}^{ij} \bar{H}_a \mathbf{10}_i^{ab} \frac{(\Sigma\Sigma\Sigma\Sigma)_b^c}{M_{\text{Pl}}^4} \mathbf{5}_c^* \\
&\quad + h_{23}^{ij} \bar{H}_a \frac{(\Sigma\Sigma)_b^a}{M_{\text{Pl}}^2} \mathbf{10}_i^{bc} \frac{(\Sigma\Sigma)_c^d}{M_{\text{Pl}}^2} \mathbf{5}_d^* + h_{24}^{ij} \bar{H}_a \frac{(\Sigma\Sigma\Sigma)_b^a}{M_{\text{Pl}}^3} \mathbf{10}_i^{bc} \frac{\Sigma_c^d}{M_{\text{Pl}}} \mathbf{5}_d^* \\
&\quad \left. + h_{25}^{ij} \bar{H}_a \frac{\Sigma_b^a}{M_{\text{Pl}}} \mathbf{10}_i^{bc} \frac{(\Sigma\Sigma\Sigma)_c^d}{M_{\text{Pl}}^3} \mathbf{5}_d^* \right). \quad (6)
\end{aligned}$$

Each matrix element h^{ij} is assumed to have an $\mathcal{O}(1)$ coefficient, and the mass hierarchy is produced by the suppression factors $\sigma/M_{\text{Pl}} \equiv 1/a$ [13].

Decomposing the superpotential Eqs.(3)-(6) into its component fields, we obtain the Yukawa couplings of down-type quarks and charged leptons as

$$\begin{aligned}
Y_d &= Y' - \frac{3}{a} h'_1 + \frac{2}{a} h'_2 + \frac{9}{a^2} h'_4 + \frac{4}{a^2} h'_5 - \frac{6}{a^2} h'_6 \\
&\quad - \frac{27}{a^3} h_{10} + \frac{8}{a^3} h_{11} + \frac{18}{a^3} h_{12} - \frac{12}{a^3} h_{13} + \dots, \quad (7)
\end{aligned}$$

$$\begin{aligned}
Y_e &= Y' - \frac{3}{a} h'_1 - \frac{3}{a} h'_2 + \frac{9}{a^2} h'_4 + \frac{9}{a^2} h'_5 + \frac{9}{a^2} h'_6 \\
&\quad - \frac{27}{a^3} h_{10} - \frac{27}{a^3} h_{11} - \frac{27}{a^3} h_{12} - \frac{27}{a^3} h_{13} + \dots, \quad (8)
\end{aligned}$$

where the matrices with a prime symbol are defined as

$$Y' \equiv Y_2 + \frac{30}{a^2} h_3 - \frac{30}{a^3} h_9 + \frac{900}{a^4} h_{14} + \frac{210}{a^4} h_{15}, \quad (9)$$

$$h'_{1,2} \equiv h_{1,2} + \frac{30}{a^2} h_{7,8} - \frac{30}{a^3} h_{16,17}, \quad (10)$$

$$h'_{4,5,6} \equiv h_{4,5,6} + \frac{30}{a^2} h_{18,19,20}. \quad (11)$$

On the other hand, the couplings Y_{ql} and Y_{ud} which are associated with the interactions $Q_i \epsilon L_j \bar{H}_C$ and $u_{Ri}^c d_{Rj}^c \bar{H}_C$, respectively, are given by

$$\begin{aligned}
Y_{ql} &= Y' + \frac{2}{a} h'_1 - \frac{3}{a} h'_2 + \frac{4}{a^2} h'_4 + \frac{9}{a^2} h'_5 - \frac{6}{a^2} h'_6 \\
&\quad + \frac{8}{a^3} h_{10} - \frac{27}{a^3} h_{11} - \frac{12}{a^3} h_{12} + \frac{18}{a^3} h_{13} + \dots, \quad (12)
\end{aligned}$$

$$\begin{aligned}
Y_{ud} &= Y' + \frac{2}{a} h'_1 + \frac{2}{a} h'_2 + \frac{4}{a^2} h'_4 + \frac{4}{a^2} h'_5 + \frac{4}{a^2} h'_6 \\
&\quad + \frac{8}{a^3} h_{10} + \frac{8}{a^3} h_{11} + \frac{8}{a^3} h_{12} + \frac{8}{a^3} h_{13} + \dots. \quad (13)
\end{aligned}$$

The terms in which the adjoint Higgs fields are contracted by themselves have larger pre-factors than the others. Such terms should be practically regarded as lower order contribution. For example, we should regard h_3 as the *first order* term like h_1 and h_2 . The terms of $h_{7,9}$ and h_{14} are referred as the *second order* terms such as $h_{4,6}$, and those of h_{15-20} should belong to the *third order* terms such as h_{10-13} .

The value of σ is related to the mass of the coloured Higgs triplet M_C and the GUT scale. The GUT scale is represented as $(M_V^2 M_\Sigma)^{1/3}$, where M_V stands for mass of X and Y bosons ($SU(5)$ breaking gauge bosons) and M_Σ for the mass of Σ . The magnitudes of these mass parameters are strictly constrained from the gauge coupling unification condition[6, 11]. However, the value of σ itself can be larger than the GUT scale $(M_V^2 M_\Sigma)^{1/3} \simeq 2.0 \times 10^{16}$ GeV by taking the Higgs couplings among H , $\bar{H}$, and Σ to be small. Therefore, we can take $a = \mathcal{O}(10 \sim 100)$, where dimension six proton decay operators are suppressed enough.

Let us now illustrate a concrete example for the Yukawa couplings which reproduces not only the realistic fermion mass spectrum but also the completely vanishing dimension five proton decay operators. We take the basis where Y_d and Y_e are diagonal. The coefficients of the dimension five proton decay operators are denoted

$$C_{5L}^{ijkl} \equiv Y_{ql}^{ij} Y_{qq}^{kl}, \quad C_{5R}^{ijkl} \equiv Y_{ud}^{ij} Y_{eu}^{kl}, \quad (14)$$

where Y_{qq} and Y_{eu} are the couplings of interactions of the coloured Higgs field coming from the $\mathbf{10}_i \mathbf{10}_j H$ type terms. Avoiding unreliable large couplings in h_i 's, the

texture of h_i 's for the third generation is uniquely determined except for $\Delta y_3 \equiv y_\tau - y_b$ which is the difference between the Yukawa couplings of the tau lepton and the bottom quark. It should be small at the GUT scale and we assume that it is provided by *second order* terms. From Eqs.(7)-(13), the third generation Yukawa components are induced as

$$(h_6)_{33} = a^2 \Delta y_3 / 25, \quad (15)$$

$$(h_5)_{33} = a (h'_2)_{33} + 2a^2 \Delta y_3 / 25, \quad (16)$$

$$(h_4)_{33} = a (h'_1)_{33} + a^2 y_b / 5 + 2a^2 \Delta y_3 / 25, \quad (17)$$

$$(h'_3)_{33} = -2a^2 (y_b + \Delta y_3) / 75 - a (h'_1 + h'_2)_{33} / 5, \quad (18)$$

up to the *second order*. Here, $h'_3 \equiv h_3 - h_9/a + 30h_{14}/a^2$. In order to avoid $\mathcal{O}(a^2 y_b)$ terms in $(h_5)_{33}$ and $(h_4)_{33}$, we must take

$$(h_1)_{33} = -ay_b/5, \quad (h_2)_{33} = 0. \quad (19)$$

Then, the value of $(h_3)_{33}$ is determined up to order $\mathcal{O}(a^2 y_b)$ as

$$(h_3)_{33} = a^2 y_b / 75. \quad (20)$$

This term should be regarded as the *first order* term because of the large pre-factor $1/75$, in which the value of $(h_3)_{33}$ itself is kept of $\mathcal{O}(1)$. Now all the third generation components of h_i 's are determined. The couplings for the second generation are also determined in a similar way. Those for the first generation are not uniquely determined since there are large degrees of freedom in the *third order* terms.

Summarizing above discussions, the *first order* terms are uniquely determined as

$$h_1 = -\frac{a}{5} \text{diag}(0, 0, y_b), \quad h_2 = \mathbf{0},$$

$$h_3 = \frac{a^2}{75} \text{diag}(0, 0, y_b). \quad (21)$$

The *second order* terms are also determined almost automatically as

$$h_4 = \frac{a^2}{75} \text{diag}(0, 9y_s + y_\mu, \Delta y_3),$$

$$h_5 = \frac{a^2}{75} \text{diag}(0, -6y_s + y_\mu, \Delta y_3),$$

$$h_6 = \frac{a^2}{25} \text{diag}(0, -y_s + y_\mu, \Delta y_3),$$

$$h_7 = h_8 = -\frac{a^3}{450} \text{diag}(0, y_\mu, \Delta y_3),$$

$$h_9 = h_{14} = \mathbf{0}. \quad (22)$$

For the *third order* terms, there are various choices, and one example is

$$h_{10} = h_{16-25} = \mathbf{0},$$

$$h_{11} = \frac{a^3}{25} \text{diag}(-y_d + 2y_e/7, 0, 0),$$

$$h_{12} = \frac{a^3}{25} \text{diag}(3y_d/2 - 2y_e/3, 0, 0),$$

$$h_{13} = \frac{a^3}{25} \text{diag}(-y_d/2 - y_e/3, 0, 0),$$

$$h_{15} = \frac{4a^4}{3675} \text{diag}(y_e, 0, 0). \quad (23)$$

It is worth noting that the accurate Yukawa couplings of the down-sector quarks and charged leptons are obtained as

$$Y_d = \text{diag}(y_d, y_s, y_b), \quad Y_e = \text{diag}(y_e, y_\mu, y_\tau), \quad (24)$$

as well as the couplings of coloured Higgs triplet vanish as

$$Y_{qt} = Y_{ud} = \mathbf{0}. \quad (25)$$

Once these Yukawa interactions are realized at the GUT scale, any dimension five proton decay process will not appear even if the renormalization group equation (RGE) effects are taken into account.

	order		Example 1	Example 2
W_1	1st	h_1	(0, 0, -0.64)	
		h_2	$\mathbf{0}$	
W_2	1st	h_3	(0, 0, 2.3)	
		h_4	(0, 0.72, 0.40)	
		h_5	(0, -0.21, 0.40)	
W_3	2nd	h_6	(0, 0.29, 1.2)	
		h_7	(0, -1.5, -3.6)	
		h_8	(0, -1.5, -3.6)	
	3rd	h_9	$\mathbf{0}$	
		h_{10}	$\mathbf{0}$	$\mathbf{0}$
		h_{11}	(-0.48, 0, 0)	$\mathbf{0}$
W_4	2nd	h_{12}	(0.68, 0, 0)	$\mathbf{0}$
		h_{13}	(-0.29, 0, 0)	$\mathbf{0}$
	3rd	h_{14}	$\mathbf{0}$	
		h_{15}	(0.19, 0, 0)	(-0.65, 0, 0)
		h_{16}	$\mathbf{0}$	$\mathbf{0}$
		h_{17}	$\mathbf{0}$	$\mathbf{0}$
		h_{18}	$\mathbf{0}$	(3.3, 0, 0)
		h_{19}	$\mathbf{0}$	(-1.4, 0, 0)
		h_{20}	$\mathbf{0}$	(-0.71, 0, 0)
		h_{21}	$\mathbf{0}$	$\mathbf{0}$
		h_{22}	$\mathbf{0}$	$\mathbf{0}$
		h_{23}	$\mathbf{0}$	$\mathbf{0}$

TABLE I: Examples of the matrix elements of h_i 's at the GUT scale which reproduce the realistic fermion mass spectrum and vanish the proton decay operators. Here, we take $a = 55$ and $\tan \beta = 10$. We take the diagonal basis of the down-type quark and the charged lepton mass matrices. The flavour mixing is imposed into the up-sector Yukawa couplings.

In Table I, we present the magnitudes of Yukawa couplings at the GUT scale by using the results in Ref.[12]. The applicability of the perturbation (couplings $\lesssim \sqrt{4\pi}$) is satisfied for all components. Notice that it can be satisfied even in the rather large $\tan \beta$ ($\tan \beta \sim 10$) region. Ordinal $SU(5)$ GUT models, e.g. with decoupling SUSY breaking spectrum, should have a small $\tan \beta$ of order 1[8].

The Yukawa couplings of the up-type quarks can also be derived from higher dimensional terms. We assume

that the Yukawa coupling of the top quark comes from Y_1 in Eq.(2), that of the charm quark from W_2 , and that of the up quark from W_4 . For example, supposing the simple superpotential,

$$W = \frac{\epsilon_{abcde}}{4} \mathbf{10}_i^{ab} \mathbf{10}_j^{cd} H^e \left\{ f_c^{ij} \frac{(\Sigma\Sigma)}{M_{\text{Pl}}^2} + f_u^{ij} \frac{(\Sigma\Sigma\Sigma\Sigma)}{M_{\text{Pl}}^4} \right\},$$

we obtain the Yukawa matrices

$$Y_u = Y_1 + \frac{30}{a^2} f_c + \frac{210}{a^4} f_u.$$

In the basis Y_u should be given by $Y_u = U_{\text{CKM}}^\dagger Y_u^{\text{diag}}$. This example gives

$$(Y_1)_{33} = 0.75, \quad (f_c)_{22} = 0.18, \quad (f_u)_{11} = 0.26,$$

for Y_u^{diag} with the same values of a and $\tan\beta$ in Tab.I.

Some comments are in order. The first is about the coefficients of the proton decay operators. They must include at least one first generation quark superfield. Therefore, it is not necessary to eliminate all components of the coloured Higgs Yukawa couplings as in Eq.(25). In fact we can realize such Yukawa matrices which have more choices than the examples shown above. However, in this case the RGE effect must be taken into account to estimate the proton decay rate, since the second and third generation components of Y_{ql} and Y_{ud} are transmitted into the first generation components through the generation mixings. The RGE analysis shows that the large entry of the second and third generation components could be destructive[14]. This means that RGE effects will break the proton stability even if all the first

generation components of Y_{ql} and Y_{ud} are zero. If Y_{ql} and Y_{ud} do not include the first generation components at the nucleon mass scale (not at the GUT scale), the dimension five proton decay processes will disappear as pointed out in Ref.[7]. It is an interesting possibility. However, there must be a reason why the scale is not the GUT scale but the nucleon mass scale.

The second comment is about the contribution from the sub-leading effects. When the soft SUSY breaking tri-linear scalar interactions of the coloured Higgs

$$-\mathcal{L}_{\text{soft}} \supset A_{ql}^{ij} \tilde{Q}_i \epsilon \tilde{L}_j \bar{H}_C + A_{ud}^{ij} \tilde{u}_{Ri}^* \tilde{d}_{Rj}^* \bar{H}_C + \text{H.c.}, \quad (26)$$

are introduced, $A_{ql} Y_{qq}$ and $A_{ud} Y_{eu}$ can contribute to the proton decay processes. We have neglected these A -term contributions in the above discussions, which can be justified in the minimal supergravity context. It is because these terms are generally proportional to the corresponding Yukawa couplings. Therefore, when they do not exist at the GUT scale, there will be no contribution at the low energy scale even if we take account into the RGE effects.

We have tried to reproduce the suitable fermion mass hierarchy as well as to suppress the proton decay in the $SU(5)$ SUSY GUT framework with the minimal field contents. The realistic fermion mass spectrum can be realized simultaneously with vanishing dimension five proton decay processes. These requirements have almost determined the Yukawa couplings of higher dimensional operators at the GUT scale.

The authors would like to thank N. Okada and M. Kakizaki for useful discussions and be grateful to A. Merle for reading the manuscript carefully. N.H. is supported by Alexander von Humboldt Foundation.

-
- [1] H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32** (1974) 438.
 - [2] K. Inoue, A. Kakuto, H. Komatsu, and S. Takeshita Prog. Theor. Phys. **68** (1982) 927 [Erratum-ibid. **70** (1983) 330]; L. E. Ibanez, and G. G. Ross, Phys. Lett. **B110** (1982) 215; J. R. Ellis, D. V. Nanopoulos, and K. Tamvakis, Phys. Lett. **B121** (1983) 123; L. Alvarez-Gaume, J. Polchinski, and M. B. Wise, Nucl. Phys. **B221** (1983) 495.
 - [3] N. Sakai and T. Yanagida, Nucl. Phys. **197** (1982) 533, S. Weinberg, Phys. Rev. **D26** (1982) 287.
 - [4] Super-Kamiokande Collaboration (K. Kobayashi *et al.*), Phys. Rev. **D72** (2005) 052007.
 - [5] T. Goto and T. Nihei, Phys. Rev. **D59** (1999) 115009.
 - [6] H. Murayama and A. Pierce, Phys. Rev. **D65** (2002) 055009.
 - [7] B. Bajc, P. Fileviez Pérez and G. Senjanović, Phys. Rev. **66** (2002) 075005.
 - [8] D. Emmanuel-Costa and S. Wiesenfeldt, Nucl. Phys. **B661** (2003) 62.
 - [9] P. Nath, Phys. Rev. Lett. **76** (1996) 2218; P. Nath, Phys. Lett. **B381** (1996) 147.
 - [10] Z. Berezhiani, Z. Tavartkiladze, and M. Vysotsky, arXiv:hep-ph/9809301.
 - [11] J. Hisano, H. Murayama, and T. Yanagida, Phys. Rev. Lett. **69** (1992) 1014; J. Hisano, H. Murayama, and T. Yanagida, Nucl. Phys. **B402** (1993) 46.
 - [12] H. Fusaoka and Y. Koide, Phys. Rev. **D57** (1998) 3986.
 - [13] The authors of Ref.[8] have also introduced higher dimensional terms but up to the first order of $1/a$. In this case, there is an inevitable relation between the Yukawa matrices as $Y_{ql} - Y_{ud} = Y_e - Y_d \neq 0$, where the dimension five operators cannot completely vanish and the suppression factor $1/a$ has nothing to do with the fermion mass hierarchy. On the other hand, the author of Ref.[9] has tried to reproduce the so called Georgi-Jarlskog texture [See H. Georgi and C. Jarlskog, Phys. Lett. **B86** (1979) 297] by introducing the terms up to the third order of $1/a$.
 - [14] In Ref.[8], the authors adopted the ingenious texture which they referred as *the consistent model 1*. This model has the non-zero first generation components, however, they avoided the large RGE effects by vanishing second and third generations' components in Y_{ql} . This scenario is effective only in the case of small $\tan\beta$ such as $\tan\beta = \mathcal{O}(1)$.